

## A study of Indoor Air Quality and its Impacts on Health

SHEFALI CHOPRA\*<sup>1</sup>, PUJA GUPTA<sup>2</sup> AND MEENAKSHI MITAL<sup>3</sup>

<sup>1</sup>Assistant Professor and <sup>2&3</sup>Professor

Department of Resource Management and Design Application

Lady Irwin College, University of Delhi, New Delhi (India)

### ABSTRACT

This work fills a significant vacuum in the literature on public health by providing a thorough assessment of the state of the art on indoor air quality (IAQ) and its effects on human health. People spend a lot of time inside, yet our knowledge of the intricate connection between different indoor air contaminants and health effects is still lacking. In order to describe the main indoor air pollutants—particulate matter, volatile organic compounds, carbon monoxide, and radon—as well as their origins, health consequences, and the groups most at risk, this study synthesises data from epidemiological and toxicological investigations. Short-term health impacts such as eye irritation, headaches, and fatigue are discussed, alongside long-term consequences including respiratory diseases, cardiovascular diseases, and cancer. The paper further critiques the methodologies used in existing studies, identifying key limitations and areas for future research. Building on this foundation, a novel framework for assessing and mitigating risks associated with poor IAQ is proposed (Goyal and Khare, 2011). This framework emphasizes a multi-faceted approach encompassing source control, improved ventilation, air cleaning technologies, and the development of comprehensive indoor air quality standards. Recommendations for policy development and future research directions are provided, aiming to enhance public health outcomes through improved indoor air quality management. This review underscores the urgent need for a cohesive strategy to address the pervasive issue of indoor air pollution, calling for an integrated effort among researchers, policymakers, and practitioners in the field.

**Key Words :** Indoor air quality, Health impacts, Indoor air pollution, Respiratory health, Ventilation, Air purifiers, Regulations

### INTRODUCTION

The term “indoor air quality (IAQ) describes the quality of the air within and surrounding buildings and other structures, with a focus on the comfort and health of building inhabitants. It is possible to lower the risk of indoor health issues by being aware of and taking control of common indoor contaminants.” The importance of indoor air quality (IAQ) has grown over time, since estimates from the US Environmental Protection Agency indicate that individuals spend around 90% of their time inside (EPA). Thus, indoor spaces are very important for public health. Inside buildings, pollutants may originate from a variety of sources, such as construction materials,

furniture, electronics, cleaning supplies, and outside air pollution that seeps into interior areas (Gupta *et al.*, 2020). Poor IAQ may have short-term health implications including irritation of the eyes and throat as well as long-term repercussions like cancer, heart disease, and respiratory disorders. IAQ is a crucial field for public health research and policy creation because of the potential for large health effects.

#### **Problem Statement:**

The issue of indoor air pollution has garnered increasing attention due to its direct and indirect effects on human health. Modern buildings, with their emphasis on energy efficiency, often have reduced ventilation rates

to save energy on heating and cooling. This reduction can lead to increased concentrations of indoor air pollutants unless specific measures are taken to improve air quality. Furthermore, the wide variety of sources of indoor pollutants, combined with personal factors such as age, preexisting health conditions, and individual susceptibility, complicates the relationship between indoor air pollution and health outcomes (Kulshreshtha *et al.*, 2008). Despite the known risks, gaps in regulation, standardization, and public awareness persist, leading to inconsistencies in IAQ management across different settings. The growing body of evidence linking poor IAQ to adverse health outcomes underscores the need for a comprehensive approach to understand, monitor, and mitigate indoor air pollution.

### Objectives:

The primary goal of this paper is to provide a thorough review of the current knowledge regarding IAQ and its impact on health, with an emphasis on identifying the key pollutants, their sources, and their potential health effects. This review aims to:

1. Summarize the existing literature on indoor air pollutants and their health implications, drawing on epidemiological and toxicological studies to provide a comprehensive overview of known risks.
2. Identify gaps in the current understanding of IAQ issues, including limitations in research methodologies, inconsistencies in findings, and areas lacking in-depth exploration.
3. Propose a conceptual framework for future research that addresses these gaps and facilitates the development of more effective strategies for IAQ management.
4. Offer recommendations for policy development aimed at improving IAQ standards and guidelines, with a focus on both prevention and intervention strategies to protect public health.

### Understanding Indoor Air Quality:

#### Definition and Components:

The term “indoor air quality (IAQ) describes the quality of the air within and around buildings and other structures, with a focus on the comfort and health of the inhabitants”(Kumar, 2017). The components of IAQ encompass a variety of pollutants that can adversely affect human health, comfort, or work performance.

Common pollutants include:

- Organic compounds with a high vapour pressure at room temperature are known as volatile organic compounds, or VOCs. Paints, cleaning products, insecticides, construction supplies, and office goods like printers are examples of sources.
- A combination of liquid droplets and solid particles that are present in the air is known as particulate matter (PM). PM may be released into the atmosphere directly, as in the case of dust from building sites, or it can be created there by interactions between gases such as nitrogen oxides and sulphur dioxide. There are two types of PM:
  - PM2.5: PM2.5, or small particles with a diameter of 2.5 micrometres or less, can penetrate the respiratory tract, posing significant health risks, including respiratory and cardiovascular problems, as they bypass the nose and throat.
  - PM10: PM10, or “coarse particles” with a diameter of 10 micrometres or less, include dust, pollen, and Mold. They can irritate eyes, nose, throat, and cause heart and lung diseases.
- Carbon Monoxide (CO): A colourless, odorless gas produced by burning carbon-containing materials. Insufficient ventilation can lead to dangerous levels of CO indoors, especially from sources like heating systems and kitchen appliances.
- Radon: A radioactive gas that occurs naturally from the decay of uranium in the soil beneath buildings. Radon can enter buildings through cracks and openings in the foundation and become trapped indoors.

**Sources:** The sources of indoor air pollutants are varied and can be broadly categorized into four main groups:

- Outdoor Air Pollution Infiltration: Outdoor pollutants can enter buildings through open windows, doors, and ventilation systems. Traffic-related pollution and industrial emissions are significant sources.
- Building Materials: Many building materials and furnishings release pollutants continuously. Examples include insulation, pressed wood products, carpets, and furniture that may release formaldehyde, a potent VOC.

- Human Activities: Activities such as cooking, heating, smoking, and using cleaning products can introduce pollutants like CO, PM, and VOCs into the indoor environment.
- Biological Agents: Microbes, pet dander, pollen, and pests can contribute to poor IAQ. Moisture build-up can further exacerbate the proliferation of Mold and mildew, leading to increased allergens and microbes in the air.

### Measurement and Standards:

Assessing IAQ involves measuring the concentrations of specific pollutants and evaluating the effectiveness of ventilation systems. Methods for assessing IAQ include:

- Direct Monitoring: Using sensors and instruments to measure pollutant levels in real-time. Technologies vary from simple carbon monoxide detectors to sophisticated systems capable of detecting and quantifying a wide range of chemical and biological pollutants.
- Indirect Monitoring: Evaluating factors that influence IAQ, such as ventilation performance, humidity levels, and the presence of sources of pollutants.

### Health Impacts of Indoor Air Pollution:

Indoor air pollution is a silent threat that can have profound effects on human health, ranging from immediate, short-term problems to severe, long-term illnesses. The nature and severity of these health impacts depend on several factors, including the type and concentration of pollutants, exposure duration, and individual susceptibility.

### Sick Building Syndrome (SBS):

Sick Building Syndrome (SBS) refers to a condition where building occupants experience acute health discomfort that improves upon leaving the building, without a specific illness being identified. Symptoms include headache, eye, nose, and throat irritation, fatigue, dizziness, and sensitivity to odors. SBS is associated with factors like poor air quality, inadequate ventilation, presence of volatile organic compounds (VOCs), and particulate matter. Addressing SBS involves improving ventilation, removing pollutant sources, maintaining proper indoor humidity, and ensuring clean indoor environments. Effective management also includes ergonomic

adjustments and addressing occupants' psychosocial concerns.

### Short-term Effects:

The immediate health effects of exposure to indoor air pollutants are often acute and reversible upon reducing or eliminating exposure. Common symptoms include irritation of the eyes, nose, and throat, which can be particularly noticeable in environments with high levels of volatile organic compounds (VOCs) or particulate matter (PM). Headaches, dizziness, and fatigue are also frequent complaints, potentially linked to elevated carbon monoxide (CO) levels or the presence of other toxic substances in the air. Furthermore, dust mites, pet dander, mould spores, or pollen inside might cause allergic responses or asthma exacerbations in some people (Kumar *et al.*, 2016). Although these symptoms might make everyday tasks more difficult and lower one's quality of life overall, they usually go away as the air quality improves.


### Long-term Effects:

Prolonged exposure to low indoor air quality has more serious and perhaps fatal long-term health effects. A prolonged exposure to some pollutants, such as radon, asbestos, and fine particulate matter, may significantly increase the chance of developing respiratory disorders, such as lung cancer and chronic obstructive pulmonary disease (COPD). Because prolonged indoor air pollution exposure may worsen systemic inflammation and vascular dysfunction, it has also been related to heart disease. Moreover, long-term exposure to certain air pollutants may lead to the development of cancer, primarily in the lungs but perhaps in other organs as well, depending on the specific chemicals involved. It's crucial to address IAQ issues early on since these chronic diseases often show no symptoms for years or even decades after exposure.

### Sensitive Groups:

Certain populations are particularly vulnerable to the adverse effects of indoor air pollution due to physiological, developmental, or health-related factors. Children are at increased risk because they breathe more rapidly than adults, inhaling more pollutants relative to their body weight. Their developing organs and immune systems also make them more susceptible to the toxic effects of pollutants (Sharma *et al.*, 2020). The elderly, another

sensitive group, may have diminished physiological reserves and often live with chronic health conditions that can be exacerbated by poor air quality. Individuals with preexisting respiratory or cardiovascular diseases, as well as those with compromised immune systems, face heightened risks from indoor air pollutants. These sensitive groups require special consideration in efforts to monitor and improve IAQ, underscoring the need for stringent standards and proactive measures to protect the most vulnerable members of our communities



### Review of Literature:

(Buonanno *et al.*, 2019) “Particle emission factors during cooking activities” This study will investigate the release of airborne particles (such as PM<sub>2.5</sub> and ultrafine particles) from cooking processes and will evaluate the factors that influence emission rates. These factors include the temperature at which food is cooked, the method of cooking, the type of food, and the ventilation in the kitchen.

(Paleologos *et al.*, 2021) This paper explores indoor air pollutants, their origins, sampling, measurement, and impact on human health. It emphasises the significance of indoor environmental quality (IEQ) and its effects on comfort and health as well as the management, technology, and recommendations for indoor air quality (IAQ). The notion of environmentally-friendly buildings is examined, accompanied by the need of maximising ventilation, air conditioning, and indoor air quality systems. The significance of productivity, psychological, and health elements in building design, construction, and operation is emphasised in the study.

(Isabella Annesi *et al.*, 2013) For a school to be secure, healthy, and productive, the indoor air quality is

essential. Research indicates that increased levels of air pollution may have an effect on children’s respiratory health, school attendance, and academic achievement. People with allergies are more vulnerable to respiratory health issues. Enhancing the quality of the air is crucial for prevention.

(Cheek *et al.*, 2021), In comparison to the control, there were decreases in indoor air quality (PM<sub>2.5</sub>) of 22.6–92.0%, according to a comprehensive assessment of the literature on portable air purifiers (PAPs). Studies have discovered correlations with outcomes related to pregnancy, respiratory parameters, blood pressure, and other health effects. Nevertheless, there was few and contradictory data to support these correlations.

Kumar *et al.* (2023). Although the importance of indoor air quality (IAQ) for health and wellbeing cannot be overstated, little is known about its consequences. Significant indoor air pollution has an impact on women’s, foetal, neurological, ophthalmic, and cardiovascular health in addition to degrading indoor air quality. Because of their physical attributes and longer exposure periods, women and children are more vulnerable. IAQ exposure may cause low birth weight, respiratory illnesses, and premature mortality, among other health problems. Reducing health dangers requires addressing research requirements and putting effective treatments into place.

Holden *et al.*, (2023) Poor housing quality significantly impacts indoor air quality, affecting respiratory health in children and young people. Factors like dampness, mold, cold homes, and pests contribute to poor indoor air quality. Children from lower-income backgrounds are at risk. Research shows that rehousing or modifying poor-quality housing can improve indoor air quality. Urgent action is needed to address this risk factor and work with families.

Saini *et al.* (2020), In poor nations, indoor air pollution (IAP) is a serious issue that has an impact on morbidity and death rates. Coal and biomass are used by around 3 billion people for home energy, which has an impact on productivity and well-being. This article presents new concepts and avenues for IAQ monitoring, along with a discussion of wireless technologies for microcontrollers and real-time monitoring.

(Mannan and Al-Ghamdi, 2021), Since most individuals spend 90% of their time inside, pollutants and toxins from cooking, air conditioning, heating, and construction materials are present. The productivity and health of people are greatly impacted by this. In spite of

twenty years of study on indoor air quality (IAQ), a thorough assessment of peer-reviewed papers remains lacking. By identifying contaminants in indoor environments—both residential and commercial—this assessment seeks to fill knowledge gaps and highlight trends in the field while also setting guidelines for environmentally friendly and sustainable construction practises.

### **Suggested Framework for Addressing Indoor Air Quality Issues:**

Addressing indoor air quality (IAQ) issues requires a comprehensive and multi-disciplinary approach. This framework outlines strategies for assessing risks, implementing mitigation measures, developing policies, and guiding future research to improve IAQ and protect public health (Singh *et al.*, 2020).

**Risk Assessment Model:** A robust model for assessing exposure risks associated with various indoor pollutants is essential for effective IAQ management. This model should:

1. Identify Pollutants: Catalog common indoor air pollutants, including chemical (e.g., VOCs, CO, formaldehyde), biological (e.g., mold, pollen), and particulate (e.g., PM2.5, asbestos) contaminants.
2. Determine Sources: Analyze potential sources of pollutants within indoor environments, including building materials, furnishings, human activities, and outdoor air infiltration.
3. Assess Exposure: Evaluate the frequency, duration, and intensity of occupant exposure to pollutants, considering building occupancy patterns and individual susceptibility.
4. Evaluate Health Risks: Use toxicological data and epidemiological studies to estimate the health risks associated with exposure levels, factoring in specific vulnerabilities of sensitive populations.

**Mitigation Framework:** Mitigating poor IAQ involves a systematic approach that includes source control, ventilation improvements, and air cleaning technologies:

1. Source Control: The most effective way to improve IAQ is to eliminate or reduce emissions from indoor sources. This can involve selecting low-emission materials, prohibiting smoking, and minimizing the use of VOC-emitting products.

2. Ventilation Improvements: Increase the introduction of outdoor air to dilute indoor pollutants. This can be achieved through natural ventilation (e.g., opening windows), mechanical ventilation (e.g., HVAC systems with effective filtration), or a combination of both.
3. Air Cleaning Technologies: Use air purifiers and cleaners to remove pollutants from indoor air. Technologies vary in effectiveness based on the types of pollutants they target (e.g., HEPA filters for particulates, activated carbon filters for gases and odors).

### **Policy and Regulation Recommendations:**

Effective policies and regulations are critical for maintaining and improving IAQ:

1. Set IAQ Standards: Develop and enforce standards for acceptable levels of key indoor pollutants, based on the latest scientific evidence and health impact assessments.
2. Building Codes and Guidelines: Implement guidelines for new construction and renovations that promote healthy indoor environments, including requirements for materials, ventilation systems, and building design.
3. Monitoring and Compliance: Establish protocols for regular IAQ monitoring in public and commercial buildings, along with mechanisms for enforcement and compliance.

### **Future Research Directions:**

Identifying gaps in current knowledge and suggesting areas for future investigation will drive improvements in IAQ management:

1. Long-term Health Studies: Conduct longitudinal studies to better understand the long-term health impacts of exposure to various indoor pollutants, especially in vulnerable populations.
2. Technological Innovations: Research and develop more effective IAQ improvement technologies, including advanced materials, building designs, and air cleaning systems.
3. Behavioral and Socioeconomic Factors: Explore the influence of occupant behavior, socioeconomic factors, and environmental justice issues on IAQ and health outcomes.

## Conclusion:

This paper has underscored the critical importance of indoor air quality (IAQ) to public health, illustrating how the air within our homes, workplaces, and public buildings can significantly impact well-being. Through a detailed examination of IAQ components, sources of indoor pollutants, their health effects, and strategies for mitigation, this review highlights the complex nature of indoor air pollution and its potential to harm human health.

Poor IAQ may cause both short-term and long-term health impacts, such as headaches, eye irritation, cancer, and chronic respiratory disorders. These effects highlight the critical need for efficient management and improvement solutions. Because some groups are more susceptible than others—children, the elderly, and those with previous medical disorders, for example—improving IAQ is not just a question of environmental health but also of equality and social justice.

Our proposed framework for addressing IAQ issues offers a comprehensive approach that includes a robust risk assessment model, systematic mitigation strategies, and clear recommendations for policy and regulation. By identifying pollutants, assessing exposure, and implementing both preventative and corrective measures, we can significantly reduce the health burdens associated with indoor air pollution.

Furthermore, the call for future research directions opens pathways for innovation in IAQ improvement technologies and deeper understanding of the long-term health impacts of exposure to indoor pollutants. Such research is crucial for developing more effective strategies and interventions.

In conclusion, the significance of IAQ to public health cannot be overstated. Comprehensive strategies that encompass risk assessment, mitigation, policy development, and research are essential to address IAQ issues effectively. By prioritizing IAQ, we can protect health, enhance quality of life, and ensure that our indoor environments are safe and supportive of overall well-being.

## REFERENCES

- Buonanno, Giorgio, Morawska, Lidia and Stabile, Luca (2019). "Particle emission factors during cooking activities". Environment International.
- Cheek, E., Guercio, V., Shrubsole, C. and Dimitroulopoulou, S. (2021). Portable air purification: Review of impacts on indoor air quality and health. *Science of the Total Environment*, **766**, 142585.
- Gupta, A., Goyal, R., Kulshreshtha, P. and Jain, A. (2020). Environmental Monitoring of PM<sub>2.5</sub> and CO<sub>2</sub> in Indoor Office Spaces of Delhi, India. In A. Sharma, R. Goyal, & R. Mittal (Eds.), *Indoor Environmental Quality* (Vol. 60, pp. 67–76). Springer Singapore. [https://doi.org/10.1007/978-981-15-1334-3\\_8](https://doi.org/10.1007/978-981-15-1334-3_8).
- Goyal, R. and Khare, M. (2011). Indoor air quality modeling for PM<sub>10</sub>, PM<sub>2.5</sub>, and PM<sub>1.0</sub> in naturally ventilated classrooms of an urban Indian school building. *Environmental Monitoring & Assessment*, **176**(1–4) : 501–516. <https://doi.org/10.1007/s10661-010-1600-7>
- Holden, K.A., Lee, A.R., Hawcutt, D.B. and Sinha, I.P. (2023). The impact of poor housing and indoor air quality on respiratory health in children. *Breathe (Sheff)*, **19**(2):230058
- Isabella Annesi-Maesano, Nour Baiz, Soutrik Banerjee, Peter Rudnai, Solenne Rive and on behalf of the SINPHONIE Group (2013). Indoor Air Quality and Sources in Schools and Related Health Effects. *J. Toxicology & Environmental Health, Part B*, **16** (8) : 491-550, DOI: 10.1080/10937404.2013.853609
- Kumar, P., Singh, A. B., Arora, T., Singh, S. and Singh, R. (2023). Critical review on emerging health effects associated with the indoor air quality and its sustainable management. *Science of The Total Environment*, **872** : 162163.
- Kulshreshtha, P., Khare, M. and Seetharaman, P. (2008). Indoor air quality assessment in and around urban slums of Delhi city, India. *Indoor Air*, **18**(6) : 488–498. <https://doi.org/10.1111/j.1600-0668.2008.00550.x>
- Kumar, P. (Ed.). (2017). Airborne particles: Origin, emissions and health impacts. Nova Science Publisher's, Inc.
- Kumar, P., Martani, C., Morawska, L., Norford, L., Choudhary, R., Bell, M. and Leach, M. (2016). Indoor air quality and energy management through real-time sensing in commercial buildings. *Energy & Buildings*, **111** : 145–153. <https://doi.org/10.1016/j.enbuild.2015.11.037>.
- Mannan, M. and Al-Ghamdi, S.G. (2021). Indoor Air Quality in Buildings: A Comprehensive Review on the Factors Influencing Air Pollution in Residential and Commercial Structure. *Internat. J. Environ. Res. Public Health*, **18**, 3276. <https://doi.org/10.3390/ijerph18063276>
- Paleologos, K.E., Selim, M.Y. and Mohamed, A.M.O. (2021). Indoor air quality: pollutants, health effects, and regulations. In *Pollution assessment for sustainable practices in applied sciences and engineering* (pp. 405-489). Butterworth-Heinemann.

Saini, J., Dutta, M. and Marques, G.A. (2020). comprehensive review on indoor air quality monitoring systems for enhanced public health. *Sustain. Environ. Res.*, **30** (6). <https://doi.org/10.1186/s42834-020-0047-y>

Sharma, A., Goyal, R. and Mittal, R. (Eds.). (2020). Indoor Environmental Quality: Select Proceedings of the 1st ACIEQ (Vol. 60). Springer Singapore. <https://doi.org/10.1007/978-981-15-1334-3>

Singh, P., Arora, R. and Goyal, R. (2020). Classroom Ventilation and Its Impact on Concentration and Performance of Students: Evidences from Air-Conditioned and Naturally Ventilated Schools of Delhi. In A. Sharma, R. Goyal, & R. Mittal (Eds.), *Indoor Environmental Quality* (Vol. **60**, pp. 125–137). Springer Singapore. [https://doi.org/10.1007/978-981-15-1334-3\\_14](https://doi.org/10.1007/978-981-15-1334-3_14)

\*\*\*\*\*